

## Publications

1. [Burns JR, Lambert GS, Baldwin MR.](#) 2017. Insights into the Mechanisms by Which Clostridial Neurotoxins Discriminate between Gangliosides. *Biochemistry*. 2017;56(20):2571-83. doi: 10.1021/acs.biochem.6b01246. PubMed PMID: 28441004.
2. [Lea WA, O'Neil PT, Machen AJ, Naik S, Chaudhri T, McGinn-Straub W, Tischer A, Auton MT, Burns JR, Baldwin MR, Khar KR, Karanicolas J, Fisher MT.](#) 2016. Chaperonin-Based Biolayer Interferometry To Assess the Kinetic Stability of Metastable, Aggregation-Prone Proteins. *Biochemistry*. 2016;55(35):4885-908. doi: 10.1021/acs.biochem.6b00293. PubMed PMID: 27505032.
3. [Lambert GS, Baldwin MR.](#) 2016. Evidence for dual receptor-binding sites in *Clostridium difficile* toxin A. *FEBS letters*. 2016;590(24):4550-63. doi: 10.1002/1873-3468.12487. PubMed PMID: 27861794.
4. [Manse JS, Baldwin MR.](#) 2015. Binding and entry of *Clostridium difficile* toxin B is mediated by multiple domains. *FEBS letters*. 2015;589(24 Pt B):3945-51. doi: 10.1016/j.febslet.2015.11.017. PubMed PMID: 26602083.
5. [Burns JR, Baldwin MR.](#) 2014. Tetanus neurotoxin utilizes two sequential membrane interactions for channel formation. *The Journal of biological chemistry*. 2014;289(32):22450-8. doi: 10.1074/jbc.M114.559302. PubMed PMID: 24973217; PMCID: PMC4139251.
6. [Karalewitz, AP, Fu, Z, Baldwin, MR, Kim, JJ and JT Barbieri.](#) 2012. Botulinum neurotoxin serotype C associates with dual ganglioside receptors to facilitate cell entry. Nov 23;287(48):40806-16. doi: 10.1074/jbc.M112.404244. Epub 2012 Oct 1.
7. [Clark, G.F., Grassi, P., Pang, P.C., Panico, M., Lafrenz, D., Drobis, E.Z., Baldwin, M.R., Morris, H.R., Haslam, S.M., Schedin-Weiss, S., Sun, W., and A. Dell.](#) 2012. Tumor biomarker glycoproteins in the seminal plasma of healthy human males are endogenous ligands for DC-SIGN. *Mol Cell Proteomics* 2012 Jan;11(1):M111.008730. Epub 2011 Oct 10
8. [Benson, M.A., Fu, Z., Kim, J.J., and M.R. Baldwin.](#) 2011. Unique ganglioside recognition strategies for clostridial neurotoxins. *J Biol Chem* 286(39):34015-22. Epub 2011 Aug. 17.
9. [Kroken, A.R., Karalewitz, A.P., Fu, Z., Baldwin, M.R., Kim, J.J., and J.T. Barbieri.](#) 2011. Unique ganglioside binding by botulinum neurotoxins C and D-SA. *FEBS J*, 278(23):4486-96. Epub 2011 May 31. Review.
10. [Karalewitz, A.P., Kroken, A.R., Fu, Z., Baldwin, M.R., Kim, J.J., and J.T. Barbieri.](#) 2010. Identification of a unique ganglioside binding loop within botulinum neurotoxins C and D-SA. *Biochemistry* 49(37):8117-26.
11. [Henkel, J.S., Baldwin, M.R. and J.T. Barbieri.](#) 2010. Toxins from bacteria. EXS 100:1-29. Review.
12. [Chen, C., Fu, Z., Kim, J.J., Barbieri, J.T., and M.R. Baldwin.](#) 2009. Gangliosides as high affinity receptors for tetanus neurotoxin. *J Biol Chem* 284(39):26569-77. Epub 2009 Jul 14.
13. [Fu, Z. Chen, C., Barbieri, J.T. Kim, J.J. and M.R. Baldwin, 2009.](#) Glycosylated SV2 and gangliosides as dual receptors for botulinum neurotoxin serotype F. *Biochemistry*, 48(24):5631-41.

14. [Baldwin, M.R. and J.T. Barbieri](#). 2009. Association of botulinum neurotoxins with synaptic vesicle protein complexes. *Toxicon* 54(5):570-4. Epub 2009 Apr 9. Review.
15. [Chen, C., Baldwin, M.R., and J.T. Barbieri](#). 2008. Molecular basis for tetanus toxin coreceptor interactions. *Biochemistry* 47(27):7179-86. Epub 2008 Jun 11.
16. [Baldwin, M.R., Tepp, W.H., Przedpelski, A., Pier, C.L., Bradshaw, M., Johnson, E.A., and J.T. Barbieri](#). 2008. Subunit vaccine against the seven serotypes of botulism. *Infect Immun* 76(3):1314-8.
17. [Pier, C.L., Pier, C.L., Tepp, W.H., Bradshaw, M., Johnson, E.A., Barbieri, J.T., and M.R. Baldwin](#). 2008. Recombinant holotoxinoid vaccine against botulism. *Infect Immun* 76(1):437-42.
18. [Baldwin, M.R. and J.T. Barbieri](#), 2007. Association of botulinum neurotoxin serotypes A and B with synaptic vesicle protein complexes. *Biochemistry*, 46(11):3200-10.
19. [Eubanks, L.M., Hixon, M.S., Jin, W., Hong, S., Clancy, C.M., Tepp, W.H., Baldwin, M.R., Malizio, C.J., Goodnough, M.C., Barbieri, J.T., Johnson, E.A., Boger, D.L., Dickerson, T.J., and K.D. Janda](#), 2007. An in vitro and in vivo disconnect uncovered through high-throughput identification of botulinum neurotoxin A antagonists. *Proc Natl Acad Sci U S A*, 104(8):2602-7.
20. [Baldwin, M.R., J.J. Kim, and J.T. Barbieri](#), 2007. Botulinum neurotoxin B-host receptor recognition: it takes two receptors to tango. *Nat Struct Mol Biol*, 14(1):9-10.